


We claim:


1. A process for preparing a chiral, nonracemic compound of the formula I

5



wherein:

- 10 R^1 is



wherein:

- 15 ring A is phenyl, an 8-, 9-, 10-, 11-, 12-, 13-, or 14-membered aromatic ring, (C₃-C₈)-cycloalkyl or a 5-, 6-, 7-, 8-, 9-, 10-, 11- or 12-membered heteroaromatic ring optionally containing one to four heteroatoms selected from the group of N, O and S;

- 20 R^3 is H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyl or phenyl;

- 25 R^4 , R^5 are each independently H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂H, OCF₂-CF₃, OCF₂-CHF₂, SCF₃, O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

n is 1, 2 or 3; and

- 30 R^2 is (C₁-C₈)-alkyl wherein one or more CH₂ groups in said (C₁-C₈)-alkyl group is optionally replaced by O, CO, S, SO or SO₂, and said (C₁-C₈)-alkyl group is optionally mono-, di-, or trisubstituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHBoc, NH-CO-C(CH₃)₃, hydroxyl,

OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, tetrazole, thiazolidin-2,4-dione, indole or (C₆-C₁₀)-aryl,

5 wherein said thiazolidin-2,4-dione and aryl groups are optionally substituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHTs, NHBoc, NHCbz, NH-CO-C(CH₃)₃, hydroxyl, OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl or tetrazole, or


10 R² is an OH protecting group (PG) selected from the group comprising benzyloxymethyl, benzyl, para-methoxybenzyl or tert-butyldimethylsilyl;

which comprises:

15 A)

a) alkylation (alk-R₂/alk-PG)

reacting cis-1,3-cyclohexanediol of the formula (II)




with a compound of the formula (III)



where R² is as defined above and

X¹ is Cl, Br, I, OMs, OTs, OTf;


30 in the presence of bases in a suitable solvent to give a racemic compound of the formula (IV)



where R^2 is as defined above;

b1) enzymatic ester formation (EF) + separation (S)

subjecting the resulting compounds of the formula (IV) to stereoselective enzymatic ester formation (EF), in which the alcohols are admixed with an acyl donor and the enzyme in an organic solvent and the resulting mixture is stirred at -20 to 80°C and, after the reaction has ended, one stereoisomer is present as an ester of the formula (V)



wherein

R^6 is $C(=O)-(C_1-C_{16})$ -alkyl, $C(=O)-(C_2-C_{16})$ -alkenyl, $C(=O)-(C_3-C_{16})$ -alkynyl, $C(=O)-(C_3-C_{16})$ -cycloalkyl,

wherein one or more carbon atoms in said $C(=O)-(C_1-C_{16})$ -alkyl, $C(=O)-(C_2-C_{16})$ -alkenyl, $C(=O)-(C_3-C_{16})$ -alkynyl and $C(=O)-(C_3-C_{16})$ -cycloalkyl groups are optionally replaced by oxygen atoms, and wherein said $C(=O)-(C_1-C_{16})$ -alkyl, $C(=O)-(C_2-C_{16})$ -alkenyl, $C(=O)-(C_3-C_{16})$ -alkynyl and $C(=O)-(C_3-C_{16})$ -cycloalkyl groups are optionally substituted by 1, 2 or 3 substituents selected from the group of F, Cl, Br, CF_3 , CN, NO_2 , hydroxyl, methoxy, ethoxy, phenyl and $CO-O(C_1-C_4)$ -alkyl or $CO-O(C_2-C_4)$ -alkenyl,

wherein said $CO-O(C_1-C_4)$ -alkyl and $CO-O(C_2-C_4)$ -alkenyl substituents are optionally substituted by 1, 2 or 3 substituents selected from the group of F, Cl, Br or CF_3 , and

R^2 is as defined above,

and the other stereoisomer is present unchanged as the alcohol of the formula (IV), and are therefore separated from each other by utilizing their different chemical or physicochemical properties (separation S)

or

5

10




20

25

30

reacting further with a compound of the formula (VI)



wherein


5 ring A, R³, R⁴, R⁵ and n are each as defined above and

X² is Cl, Br, I, OTs, Oms or OTf;

10 in the presence of bases in a suitable solvent to give the compound of the formula (I), and


e) detachment of the protecting group PG (detPG)

15 if R² is an OH protecting group (PG) as defined above under R², converting the compound of the formula (Ia)



20 wherein R¹ and PG are each as defined above,

by detaching the protecting group by known methods to a compound of the formula (VII)



25

wherein R¹ is as defined above,

f) alkylation (alk- R²)

30 then reacting it with a compound of the formula (III)




wherein X^1 and R^2 are each as defined above,

- 5 in the presence of bases in a suitable solvent to give a compound of the formula (I), the product or the enantiomeric form,


it being also possible to change the sequence of individual reaction steps as described above under A):

10




to:

15




or

20



or

25



2. The process of Claim 1, wherein the processes C) and D) are employed.

30

3. The process of Claim 2, wherein compounds of the formula (III)




35

are used wherein

X^1 is Cl, Br, I, OMs or OTs.


4. The process of Claim 3, wherein compounds of the formula (III)



- 5 are used wherein

X^1 is Cl, Br or I.


5. The process of Claim 4, wherein a compound of the formula (I)



10

is prepared where:

R^1 is



15

wherein

20 ring A is phenyl, (C₃-C₈)-cycloalkyl, a fused/bicyclic 8-, 9-, 10-, 11-, 12-, 13- or 14-membered aromatic ring, or a 5-, 6-, 7-, 8-, 9-, 10-, 11- or 12-membered heteroaromatic ring optionally containing one or more heteroatoms selected from the group of N, O and S;

25 R^3 is H, CF₃, (C₁-C₆)-alkyl, (C₃-C₈)-cycloalkyl or phenyl;

25

R^4 , R^5 are each independently H, F, Br, CF₃, OCF₃, (C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

n is 1 or 2; and


30

R^2 is (C₁-C₈)-alkyl wherein one or more CH₂ groups in the said (C₁-C₈)-alkyl group is optionally replaced by O, CO, S, SO or SO₂,

and said (C₁-C₈)-alkyl group is optionally mono-, di- or trisubstituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHBoc, NH-CO-C(CH₃)₃, hydroxyl, OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, tetrazole, thiazolidin-2,4-dione, indole or (C₆-C₁₀)-aryl,


wherein said thiazolidin-2,4-dione and (C₆-C₁₀)-aryl substituents are optionally substituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHTs, NHBoc, NHCbz, NH-CO-C(CH₃)₃, hydroxyl, OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl or tetrazole.

6. The process of Claim 5, wherein a compound of the formula (I)



is prepared wherein:

R¹ is



20

wherein

ring A is phenyl;

25 R³ is (C₁-C₄)-alkyl;

R⁴, R⁵ are each independently H, (C₁-C₄)-alkyl or O-(C₁-C₄)-alkyl;

n is 1; and

30

R² is (C₁-C₈)-alkyl wherein one or more CH₂ groups in said (C₁-C₈)-alkyl group are optionally replaced by O, CO, S, SO or SO₂, and

wherein said (C₁-C₈)-alkyl group is optionally mono-, di-, or trisubstituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHBoc, NH-CO-C(CH₃)₃, hydroxyl, OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, tetrazole, thiazolidin-2,4-dione, indole or (C₆-C₁₀)-aryl,

5

wherein said thiazolidin-2,4-dione and (C₆-C₁₀)-aryl substituents are optionally substituted by F, Cl, Br, CF₃, CN, NO₂, NHAc, NHTs, NHBoc, NHCbz, NH-CO-C(CH₃)₃, hydroxyl, OCF₃, O-(C₁-C₆)-alkyl, COOH, CO-benzyloxy, CO-O(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl or tetrazole.

10